
1633 and St., Phila. Pa



Two Cases of Fibroma of the Broad Ligament

JOHN C. DA COSTA, M.D.


Gynecologist to the Jefferson Medical Col-lege Hospital and to St. Agnes' Hospital, Philadelphia

REPRINTED FROM

THE AMERICAN JOURNAL OF OBSTETRICS Vol. XXXVIII. No. 3, 1898.

NEW YORK WILLIAM WOOD & COMPANY, PUBLISHERS 1898

> LIBRARY OCT. -12-1898



While tumors of the broad ligament arising primarily in one of the pelvic organs are by no means uncommon, it is but rarely that reference is made to tumors of the broad ligament itself existing without involvement of adjacent organs. Of the latter class I have been able to find but five or six which resemble the two cases of the kind which I present to-night.

The first of these tumors is a spherical mass six inches in diameter, removed from a patient in 1892. At the time of the operation I imagined that it was probably a fibrosarcoma, but as the woman is living and in good health six years after the removal of the growth I think that there could have been very

little if any sarcomatous tissue present.

This other tumor, the second of this kind which I have removed, is peculiar in that it is a pure fibroma and without attachment to either uterus, ovary, tube, or round ligament; no fibres could be found running to any of these organs. Among the few cases reported which resemble this was one by Boussi, originally diagnosed as a fibroid of the uterus, and which was treated by eschars placed over the sacrum and by use of morphine hypodermatically. The patient survived this treatment for five months, and an autopsy showed the presenceof a growth of the broad ligament, without other attachments, the size of an infant's head. Duncan2 reports the occurrence of bilateral myomata of the broad ligaments; Gross, of Nancy, removed a single tumor of this nature weighing 2,500 grammes; Polosson mentions a case similar to mine, but of smaller sized tumor. All these tumors seemed, according to their reporters, to possess a uterine pedicle. Praeger, of Los Angeles, reports

Read before the Section on Gynecology, College of Physicians of Philadelphia, May 19, 1898.

SURGEON GENERAL'S OFFICE

006-17-1898

two cases of fibromyomata, the first of the broad ligament, separate and distinct from the uterus, but pushing it out of its position; the second removed ten months after salpingo-oöphorectomy, and in this case a large excess of fibrous tissue was found. Cullingworth freports a case, in which the uterus and appendages were normal, from which he removed a fibroma of the broad ligament weighing 7 pounds and 5 ounces and measuring $8\frac{1}{2}x7\frac{1}{2}x6\frac{3}{4}$ inches. The growth had a short, narrow pedicle attached to the posterior surface of the broad ligament half an inch to the inner side of the ovary, and was entirely without adhesions to the uterus, ovaries, ligaments, or tubes; in this respect it resembles my case very closely.

The very best description of these tumors is given by Vautrin. He thinks they originate in the shining fibres of the broad ligament near the uterus. A tumor reported by him took its origin from the inferior internal part of the broad ligament near the uterine vessels and the lower segment of the uterus, and possessed thick, shining muscular fibres running to the anterior and posterior base of the uterus. It weighed over seven kilogrammes and had no attachment to the uterus. This writer also quotes Bushman's descriptions of cases of this kind reported by Billroth and by Rydygier; these tumors weighed 18 and 15 kilogrammes respectively. The tumor reported by Sänger and by Schmidt weighed 8 kilogrammes. Henly and Luschka take the ground that the shining fibres near the uterus spread themselves through the folds of the broad ligaments, the lower part of which contain many muscular fibres, blood vessels, and lymphatics. The tumor which I removed measured 8x7\frac{1}{2}x4\frac{3}{4} inches and was apparently entirely in the broad ligament. It had no attachments to either uterus, ovary, tube, or round ligament, and I could not discern the smallest fibre running to these organs. It had developed in the lower part of the broad ligament where the blood supply is most abundant. It had pushed up the broad ligament, and stood out in plain relief when the abdomen was opened. The growth was removed without difficulty, and it was found that the body of the tumor proper did not approach within 11 to 2 inches of the uterus, the tube and ovary being a considerable distance below. The growth was covered solely by the broad ligament, and its pedicle was composed of the broad ligament and the vessels which came from about the insertion of the uterine artery. I tied off the long, broad pedicle, measuring

4x² inches, with a cobbler's stitch, then placed a single figureof-eight ligature around it below the first ligature, leaving a stump no larger than the end of my finger. This tumor had been growing for nine years. The patient, a married woman 41 years of age, had borne her last child about twelve years ago and had not been pregnant since the growth was first noticed. Nine years ago she noticed a lump about the size of an egg in her left groin, and within the space of seven years and a half this lump had developed into a mass occupying almost the entire portion of the abdomen below the umbilicus. When I began the operation I expected to do a hysterectomy. There was one point in my mind -the limited movement of the uterus when bimanual examination was practised—which made me think that the tumor was not attached to the uterus. The reason for this limited movement was that the broad ligament enveloped both the uterus and the growth. As to the nature of the tumor, the pathologist failed to find any traces of malignancy in it, pronouncing the growth a pure fibroma. I thought it worth while to report this case as one of the few which apparently originate from those muscular fibres which abound in the lower part of the broad ligament. Almost all cases cited in the literature on the subject as tumors of the broad ligament show evidence of having originated either in the uterus, ovary, found ligament, or tube, and the great majority of such cases show evidences of malignancy. The case just recited shows neither.

/633 ARCH STREET.

REFERENCES.

1. Boussi: Soc. anat. de Paris, June, 1878, vol. liii., p. 368.

- 2. Duncan, R. B.: Australian Med. Jour., Melbourne, 1892, vol. xiv, p. 571-575
 - 3. GROSS: Congr. Français de Chirur., Paris 1892, vol vi., p. 240.
- 4. Polosson, M. A.: Soc. méd. de Lyon, 1893–94, vol. xxxiii., part 2, 91–94
- . 5. Praeger, E. Arnold: Trans Amer. Assoc. Obstet. and Gyn., 1894-95, vol. vii., pp. 481-485.
- 6. Cullingworth, C. J.: Trans. Obstet. Soc. of London, 1896, vol. xxxvii., p. 222.
 - 7. VAUTRIN: Annal. de Gynéc., Paris, 1895, vol. xliv., p. 390.
 - 8. REAMY, T. A.: Medical News, Phila., 1891, vol. lix., p. 299.
 - 9. SIBERT: La France médicale, Paris, 1897, vol. xliv., pp. 49-51.
 - 10. SMITH, R. T.: Brit. Med. Jour., London, 1889-91, vol. v., pp. 280-284,
 - 11. SUTTON, J. BLAND: Brit. Gyn. Jour., London, 1887, vol. iii., p. 494.
- 12. Treves, Frederick: Trans. Clinical Soc. London, 1892–93, vol. xxvi. pp. 101–104.

4 DA COSTA: FIBROMA OF THE BROAD LIGAMENT.

- 13. NATHAN, W. H.: Internat. Clinics, Phila., 1895, vol. iii., pp. 299-305.
- 14. LUDLAM, R.: The Clinique, Chicago, 1894, vol. xv., pp. 429-443.
- 15. Burnham, Walter: Chicago Med. Jour., 1867, vol xxiv., pp. 596-601.
 - 16. PIOGEY: Bull. Soc. anat. de Paris, 1850, vol. xxv., pp. 91-99.
 - 17. GOUGET, A.: Bull. Soc. anat. de Paris, 1892, vol. lxvii., pp. 222-224.
 - 18. GREY, C.: Brazil Med. Jour., Rio de Janeiro, No. 1, 1891.
 - 19. HALLOWELL, W. E.: AMER. JOUR. OF OBSTET., 1896, p. 64.
- 20 Holmes, Bayard: Chicago Med. Jour. and Exam., 1888, vol. lvi., pp. 54-68.
 - 21. KLETZSCH, G. A.: N. Y. Jour. Gyn. and Obstet., 1894, vol. i., p. 170.
 - 22. MALLETT, G. H.: Amer. Gyn. and Obstet. Jour., N. Y., 1891, p. 249.
 - 23. OGDEN, UZZIEL: Canadian Pract, Toronto, 1885, vol. x., p. 97.

MEDICAL JOURNALS

PUBLISHED BY

WILLIAM WOOD & COMPANY.

MEDICAL RECORD.

A WEEKLY JOURNAL OF MEDICINE AND SURGERY.

Price, \$5.00 a Year.

The Medical Record has for years been the leading organ of the medical profession in America, and has gained a world wide reputation as the recogprofession in America, and has gained a world-wide reputation as the recognized medium of intercommunication between the profession throughout the world. It is intended to be in every respect a medical newspaper, and contains among its Original Articles many of the most important contributions to nedical literature. The busy practitioner will find among the Therapeutic Hints and in the Clinical Department a large fund of practical matter, carefully condensed and exceedingly interesting. Medical News from all parts of the world is supplied through special correspondents, by mail and telegraph; New Publications and Inventions are reviewed and described; and in the Editorial Department matters of current interest are discussed in a manner which has established the Medical Record in the estimation of the whole profession as a thoroughly independent journal and the most influential publication of its class.

THE AMERICAN JOURNAL OF OBSTETRICS

AND DISEASES OF WOMEN AND CHILDREN.

Price, \$5.00 a Year (Issued Monthly).

This is not a special journal, as such are usually understood. As it gives special attention to lines which, more than any other, go to form the everyday experience of the general practitioner, its scope of usefulness is very wide.

The original articles appearing in its pages are selected with a view to the

practical value and general interest, and include many contributions from writers of wide celebrity and established reputation.

The Journal is not the organ of any society, being entirely independent. and consequently free to select for publication only such matter as will be most useful to its subscribers.

Society Proceedings, Book Reviews, and Abstracts on current literature in its scope are carefully prepared features which and to the completeness of

the Journal,

In order to add to its usefulness and attractiveness, special attention is given to the matter of illustrations, and all articles admitting of it are copiously illustrated by every available means. In fact, the Journal is presented in a form of typographical excellence unequalled by any other medical journal. A specimen copy will be sent free, if desired.

PRICES AND CLUB RATES:

Medical Record (Weekly), \$5.00 a year. American Journal of Obstetrics (Monthly), 5.00 a year. And when mailed to the same address and paid for according to our terms: Medical Record and Journal of Obstetrics, \$9.00 a year.

At the above low rates only when paid in advance to William Wood & Company or their Agents, NOT the Trade.

WILLIAM WOOD & COMPANY, 43, 45, & 47 East 10th Street, New York.

